American College of Cardiology Congress update 2020

CPD QUESTIONNAIRE

EARN FREE CPD POINTS

Complete, scan and email to **cpd@denovomedica.com**. Alternatively, you can complete this questionnaire online at **www.denovomedica.com**. Fill in your details using **clear block letters** and mark the answers with a tick (✓).

I agree that my CPD-accredited certificate will be forwarded to my e-mail address. _____
(Signature of healthcare professional)
(Online)

First Name																			Profession											
Surname																			Telephone											
HPC No.																			City											
E-mail																			Sales Rep.											

1. In the renal denervation trial of hypertensive patients not on medicine, what level of blood pressure reduction was achieved in the denervation group?

- ☐ **A** 6.7/4.5mmHg
- ☐ **B** 4.7/3.7mmHg
- ☐ **C** 3.5/3.2mmHg

2. What percentage of patients in the renal denervation group had to be treated for BP>180mmHg?

- ☐ **A** 10%
- ☐ **B** 17%
- ☐ **C** 25%

3. Wits University research in homozygous familial hypercholesterolaemia patients on high intensity statin, a PCSK-9 inhibitor and ezetimibe achieved a further LDL cholesterol reduction with evinacumab of:

- ☐ **A** 25%
- ☐ **B** 37%
- ☐ **C** 47%

4. In the Ischemia-CKD study of patients with chronic coronary syndrome and kidney injury, the invasive group undergoing cardiac catheterisation has _____ in respect of the stage of chronic kidney disease as those in the medically treated group.

- ☐ **A** Better
- ☐ **B** Poorer
- ☐ **C** The same results

5. Rivaroxaban and aspirin in the COMPASS trial of diabetic patients with stable coronary/peripheral disease resulted in a MACE reduction of what percent compared to aspirin alone?

- ☐ **A** 14%
- ☐ **B** 24%
- ☐ **C** 26%
- ☐ **D** 27%

6. In the TICO study using ultrathin bioresorbable sirolimus-eluting stents, 3 months treatment post-PCI with aspirin and ticagrelor followed by ticagrelor alone till 12 months had lower bleeding rates than 12 months of aspirin plus ticagrelor.

- ☐ **A** True
- ☐ **B** False

7. The use of colchicine post-MI in the COLCOT trial was shown to reduce costs by what percent over the trial period?

- ☐ **A** 30%
- ☐ **B** >40%
- ☐ **C** 20%

8. The DAPPA-HF trial in patients with heart failure and reduced ejection fraction treated with dapagliflozin showed an absolute reduction of CV death and hospitalisation for heart failure of:

- ☐ **A** 3%
- ☐ **B** 4%
- ☐ **C** 5%

9. A number of clinical trials now support the use of anticoagulation with P2Y12 monotherapy in patients with AF following acute coronary syndrome or PCI intervention.

- ☐ **A** True
- ☐ **B** False

10. TAVR in a real-world study in the UK in older patients with severe aortic stenosis resulted in a ____ death rate compared to conventional surgery?

- ☐ **A** Lower
- ☐ **B** The same
- ☐ **C** Higher

© 2020 deNovo Medica