

Complete, scan and email to [cpd@denovomedica.com](mailto:cpd@denovomedica.com). Alternatively, you can complete this questionnaire online at [www.denovomedica.com](http://www.denovomedica.com). Fill in your details using **clear block letters** and mark the answers with a tick (✓).

I agree that my CPD-accredited certificate will be forwarded to my e-mail address. \_\_\_\_\_

(Online)

(Signature of healthcare professional)

First Name																					Profession																				
Surname																						Telephone																			
HPC No.																						City																			
E-mail																																									

**1. The objectives of osteoporosis therapy are to:**

- A Improve bone strength
- B Reduce risk of fracture
- C Prevent rapid bone loss
- D All of the above
- E A and B
- F A and C

**2. Osteoporotic patients at very high risk for fracture include those with:**

- A A fracture within the past year
- B Multiple fractures
- C A fracture while on osteoporosis treatment
- D A fracture while on medication harmful to bone
- E All of the above
- F B, C and D

**3. Most guidelines recommend bisphosphonates as the preferred first-line treatment for patients at very high risk for fracture, followed by anabolic agents:**

- A True
- B False

**4. Under which circumstances is anabolic therapy recommended as second-line treatment for osteoporosis?**

- A New fracture on current osteoporosis treatment
- B Unacceptable rates of bone loss following compliance with appropriate antiresorptive therapy
- C Chronic glucocorticosteroid therapy
- D All of the above
- E A and B
- F B and C

**5. The anabolic effects of teriparatide result in an increase in:**

- A Total bone volume
- B Periosteal diameter
- C Cortical thickness
- D Endocortical diameter
- E All of the above
- F A, B and C
- G A, C and D

**6. From the VERO study, which statement is incorrect?**

- A Teriparatide was compared to risedronate at 24 months of treatment in postmenopausal women who had severe osteoporosis
- B Of the study cohort, >70% had prior osteoporosis therapy
- C Previous bisphosphonate therapy precludes the use of teriparatide

**7. Abaloparatide is a:**

- A Selective oestrogen receptor modulator
- B An analogue of human PTHrP
- C A sclerostin inhibitor

**8. From the ACTIVE trial, at 18 months of treatment with abaloparatide, new vertebral fractures were reduced by \_\_\_%?**

- A 65%
- B 80%
- C 86%

**9. Sclerostin is a protein that is released by the osteocytes, generally in response to oestrogen deficiency, lack of skeletal loading and glucocorticoids:**

- A True
- B False

**10. From the ARCH study, hip fractures were reduced by approximately \_\_\_% at 24 months of treatment with romosozumab?**

- A 12%
- B 20%
- C 40%