

Complete, scan and email to [cpd@denovomedica.com](mailto:cpd@denovomedica.com). Alternatively, you can complete this questionnaire online at [www.denovomedica.com](http://www.denovomedica.com). Fill in your details using **clear block letters** and mark the answers with a tick (✓).

I agree that my CPD-accredited certificate will be forwarded to my e-mail address. \_\_\_\_\_  
(Online) (Signature of healthcare professional)

First Name																				Profession																		
Surname																				Telephone																		
HPC No.																			City																			
E-mail																			Sales Rep.																			

- In South Africa, 1990-2010, cirrhotic deaths increased by:
  - A 10.4%
  - B 24%
  - C 34%
- Which blood tests are performed to diagnose hepatitis?
  - A ALT
  - B AST
  - C ALP
  - D All of the above
  - E A and B
  - F A and C
- Which of the following features of cirrhosis leads to portal hypertension?
  - A Fibrous tissue deposition
  - B Active vasoconstriction
  - C Nodule formation
  - D All of the above
- Which statement is false? In decompensated cirrhosis:
  - A Varices occur due to fluid build-up in the peritoneal cavity
  - B The kidneys respond by retaining salt and water
  - C Hepatic encephalopathy results from a build-up of toxins
- Stumbling/falling, fatigue and confusion are early-stage symptoms of:
  - A Varices
  - B Ascites
  - C Hepatic encephalopathy
- When varices manifest, two-year survival is:
  - A 30%
  - B 50%
  - C 75%
- Further to a second decompensating event, two-year survival is:
  - A 25-30%
  - B 45-50%
  - C 65-70%
- When there is kidney failure associated with liver disease, survival can be as long as five years:
  - A True
  - B False
- In the NAFLD patient with simple steatosis, what percentage shows no progressive liver disease?
  - A 10-20%
  - B 25-50%
  - C 80-90%
- What percentage of NAFLD patients develop cirrhosis within 10-20 years?
  - A 0-5%
  - B 10-20%
  - C 25-50%
- Of patients with alcoholic-associated liver disease, what percentage presents with steatohepatitis?
  - A 10-35%
  - B 50-60%
  - C 90-100%
- Which statement is true?
  - A Viral hepatitis B is now easily cured with simple oral therapy over eight weeks
  - B TDF use over five years to adequately suppress hepatitis B shows significant reduction in fibrosis
- Which of the following agents are often associated with drug-induced liver injury?
  - A Sildenafil
  - B Sibutramine
  - C NSAIDs
  - D Benzodiazepines
  - E All of the above
  - F A and B only
- Which disease of the bile duct can be well controlled with medical therapy?
  - A PBC
  - B PSC
  - C A and B
- Which of the following have established safety and efficacy to promote hepatic healing in viral, non-alcoholic and alcoholic steatohepatitis?
  - A Bovine colostrum
  - B Essential phospholipids
  - C N-acetylcysteine
  - D Milk thistle
  - E All of the above
  - F A, B and C